

## **Bibliography, [Inguinal Hernia And Strength Training](#), Jonathon Sullivan**

- Azurin DJ, Go LS, Schuricht A, et al. Endoscopic preperitoneal herniorrhaphy in professional athletes with groin pain. *J Laparoscop Adv Surg Tech* 2009; 7(1).
- Barile A, Erriquez D, Cacchio A et al. Groin pain in athletes: Role of magnetic resonance. *Radiol Med* 2000; 100(4):216-22.
- Dimitrakopoulou A, Schilders E. Sportsman's Hernia? An ambiguous term. *J Hip Preserv Surg* 2015; 3(1):16-22.
- Fitzgibbons RJ, Forse RA. Groin hernias in adults. *NEJM* 2015; 372(8):756-63.
- Flich J, Alfonso JL, Delgado F, et al. Inguinal hernia and certain risk factors. *Eur J Epidemiology* 1992; 8(2):277-282.
- Fredberg U, Kissmeyer-Nielsen P. The sportsman's hernia—fact or fiction? *Scand J Med Sci Sports* 1996; 6(4):201-204.
- Garvey JFW, Hazard H. Sports hernia or groin disruption injury? Chronic athletic groin pain: a retrospective study of 100 patients with long-term followup. *Hernia* 2013; 18:815-823.
- Kockerling F, Schug-Pass C, Adolf D, et al. Bilateral and unilateral total extraperitoneal inguinal hernia repair have equivalent early outcomes: analysis of 9395 cases. *World J Surg* 2015; 39:1887-1894.
- Jacob DA, Hackl JA, Bittner R, et al. Perioperative outcome of unilateral vs. bilateral inguinal repairs in TAPP technique: analysis of 15,176 cases from the Herniated Registry. *Surg Endoscopy* 2015; 29:3733-3740.
- Kumar A, Doran J, Nguyen-Van-Tam JS, Beckingham J. Results of inguinal canal repair in athletes with sports hernia. *J R Coll Edinburgh* 2002; 47(3):561-5.
- Le CB, Zadeh J, Ben-David K. Total extraperitoneal laparoscopic inguinal hernia repair with adductor tenotomy: a 10-year experience in the treatment of athletic pubalgia. *Surg Endoscopy* 2020; doi: 10.1007/s00464-020-07705-2.
- LeBlanc KE, LeBlanc LL, LeBlanc K. Inguinal hernias: Diagnosis and management. *Am Fam Phys* 2013; 87(12):844-848.
- Lockhart K, Dunn D, Teo S, et al. Mesh vs. non-mesh for inguinal and femoral hernia repair. *Cochrane database of systematic reviews* 2018; 9: DOI: 10.1002/14651858.CD011517.pub2.
- Orchard J, Read JW, Verrall GM, Slavotinek JP. Pathophysiology of chronic groin pain in the athlete. *Internat Sports Med J* 2000; 1(1): 1-16.
- Lal P, Phillips P, Chander J, Ramteke V. Is unilateral laparoscopic TEP inguinal hernia repair a job half done? The case for bilateral repair. *Surg Endoscopy* 2010; 24:1737-1745.
- Ruhl CE, Everhart JE. Risk factors for inguinal hernia among adults in the US population. *Am J Epidem* 2007; 165(10):1154-1161.
- Saggar VR, Sarangi R. Occult hernias and bilateral endoscopic total extraperitoneal inguinal hernia repair: is there a need for prophylactic repair. *Hernia* 2007; 11:47-49.
- Samir P, Poston GJ. It is highly unlikely that the development of an abdominal wall hernia can be attributed to a single strenuous event. *Ann R Coll Surg Engl* 2006; 88:168-71.
- Smith GD, Crosby DL, Lewis PA. Inguinal hernia and a single stressful event. *Ann R Coll Surg Engl* 1996; 78:367-68.

Svendson SW, Frost P, Vestergaard M, Anderson JH. Risk and prognosis of inguinal hernia in relation to occupational mechanical exposures—a systematic review of the epidemiological evidence. *Scan J Work Env Health* 2013; 39(1):5-26.

Zheng R, Altieri MS, Yang J, et al. Long-term incidence of contralateral primary hernia repair following unilateral hernia repair in a cohort of 32,824 patients. *Surg Endoscopy* 2017; 31:817-822.